

Table 6. Estimated numbers of AIDS cases in adult and adolescent Hispanics, by transmission category and place of birth, 2004—United States

Transmission category	Place of birth											
	United States		Central/South America		Cuba		Mexico		Puerto Rico		Total ^a	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact	1,182	43	437	52	121	59	860	60	313	20	3,642	42
Injection drug use	704	25	105	13	26	13	162	11	617	39	2,095	24
Male-to-male sexual contact and injection drug use	147	5	19	2	9	5	48	3	66	4	352	4
Heterosexual contact	703	25	264	32	47	23	354	25	588	37	2,478	29
Other ^b	32	1	11	1	1	0	18	1	11	1	96	1
Total^c	2,769	100	836	100	204	100	1,443	100	1,595	100	8,663	100

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

^a Includes 269 persons whose place of birth is not among those listed and 1,548 persons whose place of birth is unknown. Because row totals were calculated independently of the values for the subpopulations, the values in each row may not sum to the row total.

^b Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

^c Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.